arXiv:1212.0196v1 [math.NT] 2 Dec 2012

ON NON-CONGRUENT NUMBERS WITH 3 MODULO 8 PRIME FACTORS

SHENXING ZHANG

Abstract. In this paper, we will give a series of non-congruent numbers with $\equiv 3 \pmod 8$ prime factors.

1. Introduction

In this paper, we will give a series of non-congruent numbers with $\equiv 3 \pmod 8$ prime factors. Set

$$\left[\frac{a}{n}\right] := \frac{1}{2}\Big(1 - \Big(\frac{a}{n}\Big)\Big). \tag{1}$$

Then we have

Theorem 1.1. *Suppose $n = m$ or $2m, m = p_1 \cdots p_k$ and $p_i \equiv 3 \pmod 8$. Let A be the $k \times k$ matrix over $\mathbb{F}_2$ with (i, j)-entry $\left[\frac{p_j}{p_i}\right]$ $(i \neq j)$ and (i, i)-entry $\left[\frac{m/p_i}{p_i}\right]$. Assume $A^2 + A + I$ is invertible over $\mathbb{F}_2$. Then if $n = m$ odd, or $n = 2m$ even and k is even, we have*

$$\operatorname{rank} E(\mathbb{Q}) = \operatorname{rank} E'(\mathbb{Q}) = 0, \tag{2}$$

$$\begin{aligned} &\text{Ш}(E/\mathbb{Q})[2^\infty] = 0, \\ &\text{Ш}(E'/\mathbb{Q})[2^\infty] \cong \begin{cases} (\mathbb{Z}/2\mathbb{Z})^{k-2}, & \text{if } n \text{ odd, } 2 \mid k; \\ (\mathbb{Z}/2\mathbb{Z})^{k-1}, & \text{if } n \text{ odd, } 2 \nmid k; \\ (\mathbb{Z}/2\mathbb{Z})^{k}, & \text{if } n \text{ even, } 2 \mid k. \end{cases} \end{aligned} \tag{3}$$

Corollary 1.2. *Suppose $m = p_1 \cdots p_k$ and $p_i \equiv 3 \pmod 8$. If $\left(\frac{p_i}{p_j}\right) = 1$ for all $1 \leq i < j \leq k$, then m is a non-congruent number. If moreover k is even, then $2m$ is also a non-congruent number. In particular, we can construct an infinite set S of $\equiv 3 \pmod 8$ primes, such that the product of any finite primes in S is an non-congruent number.*

It's easy to construct several p's in S, for example,

$$S = \{3, 11, 83, 107, 659, 1019, 1619, \dots\}.$$

The product of any finite primes of them is a non-congruent number.

Date: December 4, 2012.

2. Review of 2-descent method.

In this section, we recall the 2-descent method of computing the Selmer groups of elliptic curves. One can find more details in [4]. Suppose n is a fixed odd positive square-free integer, $K = \mathbb{Q}$, and $E/\mathbb{Q}$, $E'/\mathbb{Q}$, φ, $\psi = \varphi^\vee$ are given by

$$E = E^{(n)} : y^2 = x^3 - n^2x, \quad E' = \widehat{E^{(n)}} : y^2 = x^3 + 4n^2x,$$

$$\varphi : E \to E',\ (x, y) \mapsto (\frac{y^2}{x^2}, \frac{y(x^2 + n^2)}{x^2}),$$

$$\psi : E' \to E,\ (x, y) \mapsto (\frac{y^2}{4x^2}, \frac{y(x^2 - 4n^2)}{8x^2}).$$

Then $\varphi\psi = [2], \psi\varphi = [2]$ and the following diagram of exact sequences commutes (cf. [7] p 5):

$$\begin{tikzcd}
 & 0 \arrow[d, dashed, "\iota_1"] & 0 \arrow[d, dashed, "\iota_2"] & 0 \arrow[d] & \\
0 \arrow[r] & E'(\mathbb{Q})/\varphi E(\mathbb{Q}) \arrow[r] \arrow[d, "\psi"] & S^{(\varphi)}(E/\mathbb{Q}) \arrow[r] \arrow[d] & \text{Ш}(E/\mathbb{Q})[\varphi] \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & E(\mathbb{Q})/\psi\varphi E(\mathbb{Q}) \arrow[r] \arrow[d] & S^{(\psi\varphi)}(E/\mathbb{Q}) \arrow[r] \arrow[d] & \text{Ш}(E/\mathbb{Q})[\psi\varphi] \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & E(\mathbb{Q})/\psi E'(\mathbb{Q}) \arrow[r] \arrow[d] & S^{(\psi)}(E'/\mathbb{Q}) \arrow[r] & \text{Ш}(E'/\mathbb{Q})[\psi] \arrow[r] & 0 \\
 & 0 & & &
\end{tikzcd}$$

In this case ι_1 and ι_2 are exact. Let $\tilde{S}^{(\psi)}(E'/\mathbb{Q})$ denote the image of $S^{(\psi\varphi)}(E/\mathbb{Q})$ in $S^{(\psi)}(E'/\mathbb{Q})$. Then

$$\#\text{Ш}(E/\mathbb{Q})[\varphi] = \frac{\#S^{(\varphi)}(E/\mathbb{Q})}{\#E'(\mathbb{Q})/\varphi E(\mathbb{Q})}, \quad \#\text{Ш}(E'/\mathbb{Q})[\psi] = \frac{\#S^{(\psi)}(E'/\mathbb{Q})}{\#E(\mathbb{Q})/\psi E'(\mathbb{Q})},$$

and

$$\#\text{Ш}(E/\mathbb{Q})[2] = \frac{\#S^{(\varphi)}(E/\mathbb{Q}) \cdot \#\tilde{S}^{(\psi)}(E'/\mathbb{Q})}{\#E'(\mathbb{Q})/\varphi E(\mathbb{Q}) \cdot \#E(\mathbb{Q})/\psi E'(\mathbb{Q})}. \tag{4}$$

Similarly, if we swap E and E', φ and ψ, then ι_1 and ι_2 is no more exact but their kernels coincide, we still have

$$\#\text{Ш}(E'/\mathbb{Q})[2] = \frac{\#S^{(\psi)}(E'/\mathbb{Q}) \cdot \#\tilde{S}^{(\varphi)}(E/\mathbb{Q})}{\#E(\mathbb{Q})/\psi E'(\mathbb{Q}) \cdot \#E'(\mathbb{Q})/\varphi E(\mathbb{Q})}. \tag{5}$$

The 2-descent method to compute the Selmer groups $S^{(\varphi)}(E/\mathbb{Q})$ and $S^{(\psi)}(E'/\mathbb{Q})$ is as follows (cf. [6] for general elliptic curves). Let

$$S = \{\text{prime factors of } 2n\} \cup \{\infty\},$$

$$\mathbb{Q}(S, 2) = \{b \in \mathbb{Q}^\times/\mathbb{Q}^{\times 2} : 2 \mid \operatorname{ord}_p(b), \forall p \notin S\} = \{d \in \mathbb{Z} : d \mid n\}.$$

By the exact sequence

$$0 \to E'(\mathbb{Q})/\varphi E(\mathbb{Q}) \xrightarrow{i} \mathbb{Q}(S, 2) \xrightarrow{j} WC(E/\mathbb{Q})[\varphi],$$

where

$$i:\ (x,y)\mapsto x,\ O\mapsto 1,\ (0,0)\mapsto 4n^2,\qquad j:d\mapsto\{C_d/\mathbb{Q}\}$$

and $C_d/\mathbb{Q}$ is the homogeneous space for $E/\mathbb{Q}$ defined by the equation

$$C_d: dw^2 = d^2 + 4n^2z^4, \tag{6}$$

the φ-Selmer group $S^{(\varphi)}(E/\mathbb{Q})$ is then

$$S^{(\varphi)}(E/\mathbb{Q}) \cong \{d\in\mathbb{Q}(S,2): C_d(\mathbb{Q}_p)\neq\emptyset,\ \forall p\in S\}. \tag{7}$$

Similarly, suppose

$$C'_d: dw^2 = d^2 - n^2z^4. \tag{8}$$

The ψ-Selmer group $S^{(\psi)}(E'/\mathbb{Q})$ is then

$$S^{(\psi)}(E'/\mathbb{Q}) \cong \{d\in\mathbb{Q}(S,2): C'_d(\mathbb{Q}_p)\neq\emptyset,\ \forall p\in S\}. \tag{9}$$

3. Local computation

In this section, p will be always assumed to be an odd prime number. As in [4], we need a modification of the Legendre symbol. For $x\in\mathbb{Q}_p$ or $\in\mathbb{Q}$ such that $\operatorname{ord}_p(x)$ is even, we set

$$\left(\frac{x}{p}\right) := \left(\frac{xp^{-\operatorname{ord}_p(x)}}{p}\right). \tag{10}$$

Thus $\left(\frac{-}{p}\right)$ defines a homomorphism from $\{x\in\mathbb{Q}^\times/\mathbb{Q}^{\times 2}:\operatorname{ord}_p(x)\text{ is even}\}$ to $\{\pm1\}$.

3.1. Computation of Selmer groups.

In this subsection, we list the conditions when C_d or C'_d is locally solvable, cf. [4], also cf. [7].

3.1.1. *The case n odd.*

Lemma 3.1. *$d\in S^{(\varphi)}(E/\mathbb{Q})$ if and only if d satisfies*

1. *$d>0$ has no prime factor $p\equiv 3 \pmod 4$;*
2. *$\left(\frac{n/d}{p}\right)=1$ for all odd $p\mid d$;*
3. *$\left(\frac{d}{p}\right)=1$ for all odd $p\mid(2n/d)$;*
4. *if $2\mid d$, $n\equiv\pm1 \pmod 8$.*

Lemma 3.2. *$d\in S^{(\psi)}(E'/\mathbb{Q})$ if and only if d satisfies*

1. *$d\equiv\pm1 \pmod 8$ or $n/d\equiv\pm1 \pmod 8$*
2. *$\left(\frac{n/d}{p}\right)=1$ for all $p\mid d, p\equiv 1 \pmod 4$;*
3. *$\left(\frac{d}{p}\right)=1$ for all $p\mid(n/d), p\equiv 1 \pmod 4$.*

3.1.2. *The case n even.*

Lemma 3.3. *$d \in S^{(\varphi)}(E/\mathbb{Q})$ if and only if d satisfies*

1. *$d > 0$ is odd;*
2. *d has no prime factor $p \equiv 3 \pmod 4$;*
3. *$\left(\dfrac{n/d}{p}\right) = 1$ for all odd $p \mid d$;*
4. *$\left(\dfrac{d}{p}\right) = 1$ for all odd $p \mid (n/d)$;*

Lemma 3.4. *$d \in S^{(\psi)}(E'/\mathbb{Q})$ if and only if d satisfies*

1. *$\left(\dfrac{n/d}{p}\right) = 1$ for all $p \mid d, p \equiv 1 \pmod 4$;*
2. *$\left(\dfrac{d}{p}\right) = 1$ for all $p \mid (n/d), p \equiv 1 \pmod 4$.*

3.2. **Computation of the images of Selmer groups.** We will use the following lemma to compute $\tilde{S}^{(\psi)}(E/\mathbb{Q})$, which follows from [1] §5, Lemma 10:

Lemma 3.5. *Let $d \in S^{(\psi)}(E'/\mathbb{Q})$. Suppose (σ, τ, μ) is a nonzero integer solution of $d\sigma^2 = d^2\tau^2 - n^2\mu^2$. Let $\mathcal{M}_b$ be the curve corresponding to $b \in \mathbb{Q}^\times/\mathbb{Q}^{\times 2}$ given by*

$$\mathcal{M}_b: \; dw^2 = d^2t^4 - n^2z^4, \quad d\sigma w - d^2\tau t^2 + n^2\mu z^2 = bu^2. \tag{11}$$

Then $d \in \tilde{S}^{\psi}(E'/\mathbb{Q})$ if and only if there exists $b \in \mathbb{Q}(S, 2)$ such that $\mathcal{M}_b$ is locally solvable everywhere.

Note that the existence of σ, τ, μ follows from Hasse-Minkowski theorem (cf. [5]).

Suppose $d \in \tilde{S}^{(\psi)}(E'/\mathbb{Q})$. Write $d = \tau^2 - \mu^2$ and select the triple (σ, τ, μ) in Lemma 3.5 to be $(nd, n\tau, d\mu)$. Then the defining equations of $\mathcal{M}_{nd^2b}$ in (11) can be written as

$$w^2 = d(t^4 - (n/d)^2z^4), \quad w - \tau t^2 + (n/d)\mu z^2 = bu^2. \tag{12}$$

By abuse of notations, we denote the above curve by $\mathcal{M}_b$. We use the notation $O(p^m)$ to denote a number with p-adic valuation $\geq m$.

The case $p \mid d$. We may assume

$$p \mid (\tau - \mu), \quad p \nmid (\tau + \mu).$$

It's easy to see $v(t) = v(z) \leq v(w)$. We may assume $z = 1, v(t) = 0, v(w) \geq 0$. It's easy to see $t^2 \equiv \pm\frac{n}{d} \pmod p$.

(i) If $v(bu^2) = m \geq 3$, then by $w^2 = (\tau t^2 - \frac{n\mu}{d} + O(p^m))^2 = d(t^4 - (\frac{n}{d})^2)$,

$$\left(\mu t^2 - \frac{n\tau}{d}\right)^2 = O(p^m).$$

Let $t^2 = \frac{n\tau}{d\mu} + \beta \equiv n/d \pmod p$, where $v(\beta) = \alpha \geq \frac{m}{2}$, then $\boxed{\left(\dfrac{n/d}{p}\right) = 1}$ and

$$\begin{aligned} w^2 =& d\left(-(\frac{n}{d})^2 + (\frac{n\tau}{d\mu})^2 + 2\frac{n\tau}{d\mu}\beta + \beta^2\right) \\ =& \frac{n^2}{\mu^2}(1 + \frac{2\tau\mu}{n}\beta + \frac{d\mu^2}{n^2}\beta^2), \end{aligned}$$

Take the square root on both sides, then

$$\begin{aligned} w &= \pm \frac{n}{\mu}\left(1 + \frac{1}{2}\left(\frac{2\tau\mu}{n}\beta + \frac{d\mu^2}{n^2}\beta^2\right) - \frac{1}{8}\left(\frac{2\tau\mu}{n}\beta\right)^2 + O(p^{3\alpha-3})\right) \\ &= \pm\left(\frac{n}{\mu} + \tau\beta - 2n\mu\left(\frac{\mu\beta}{2n}\right)^2 + O(p^{3\alpha-2})\right), \end{aligned}$$

but on the other hand,

$$w = \tau t^2 - \frac{n\mu}{d} + bu^2 = \frac{n}{\mu} + \tau\beta + bu^2.$$

The sign must be positive and

$$bu^2 = -2n\mu\left(\frac{\mu\beta}{2n}\right)^2 + O(p^{3\alpha-2}),$$

thus $\boxed{p \mid b,\ \left(\frac{n/b}{p}\right) = \left(\frac{-2\mu}{p}\right) = \left(\frac{-2\tau}{p}\right)}$.

(ii) If $v(bu^2) \le 2$ and $t^2 \equiv n/d \pmod p$, then $\boxed{\left(\frac{n/d}{p}\right) = 1}$. Let $t^2 = \frac{n}{d} + \frac{p^2\alpha}{2n}$, then

$$w^2 = d\cdot\frac{p^2\alpha}{2n}\cdot\left(\frac{2n}{d} + \frac{p^2\alpha}{2n}\right) = p^2\alpha\left(1 + \frac{p^2 d\alpha}{4n^2}\right),$$

and

$$\begin{aligned} w =& p\sqrt{\alpha}\left(1 + \frac{p^2 d\alpha}{8n^2} + O(p^2)\right) \\ bu^2 =& w - \tau t^2 + \frac{n\mu}{d} \\ =& \pm p\sqrt{\alpha}\left(1 + \frac{p^2 d\alpha}{8n^2}\right) - \frac{\tau n}{d} + \frac{n\mu}{d} - \frac{p^2\tau\alpha}{2n} + O(p^3) \\ =& -\frac{p^2\tau}{2n}\left(\sqrt{\alpha} \mp \frac{n}{p\tau}\right)^2 - \frac{n}{2\tau d}(\tau-\mu)^2 \pm \frac{p^3 d}{8n^2}\alpha^{3/2} + O(p^3). \end{aligned}$$

If $v(bu^2) = 2$, then $\sqrt{\alpha} \equiv \pm\frac{n}{p\tau} \pmod p$, and

$$\begin{aligned} bu^2 &= -\frac{n}{2\tau d}(\tau-\mu)^2 + \frac{p^3 d}{8n^2}\left(\frac{n}{p\tau}\right)^3 + O(p^3) \\ &= -\frac{n}{8\tau^3 d}\left(4\tau^2(\tau+\mu)^2 - d^2\right) + O(p^3) \\ &= -\frac{n}{8\tau^3 d}(\tau-\mu)^3(3\tau+\mu) + O(p^3) = O(p^3), \end{aligned}$$

which is impossible! Thus $v(bu^2) = 1$ and $\boxed{p \mid b,\ \left(\frac{n/b}{p}\right) = \left(\frac{-2\tau}{p}\right)}$.

(iii) If $v(bu^2) \le 2$ and $t^2 \equiv -n/d \pmod p$, then $\boxed{\left(\frac{-n/d}{p}\right) = 1}$ and

$$bu^2 = w - \tau t^2 + n\mu/d = 2n\tau/d + O(p),$$

thus $\boxed{p \nmid b,\ \left(\frac{b}{p}\right) = \left(\frac{2n\tau/d}{p}\right) = \left(\frac{-2\tau}{p}\right)}$.

The case $p \mid \frac{n}{d}$. We may assume $p \nmid \tau$.

(i) If $v(z) \geq v(t) = v(w)/2$, we may assume $t = 1, v(w) = 0, v(t) \geq 0$. Then

$$\begin{aligned} w &= \pm\sqrt{d}(1 - 2(mz^2/d)^2 - 2(mz^2/d)^4 + \cdots) \\ &= \tau - \mu \cdot 2mz^2/d + bu^2. \end{aligned}$$

Notice that $(\sqrt{d} - \tau)(-\sqrt{d} - \tau) = \mu^2$ and $\pm\sqrt{d} - \tau$ are co-prime. Choose suitable $\sqrt{d}$ or τ such that $\sqrt{d} - \tau \neq 0$, then $\mathrm{ord}_p(\sqrt{d} - \tau)$ is even and $\left(\frac{\sqrt{d}-\tau}{p}\right)$ is well defined.

We may assume that $p \nmid (\sqrt{d} + \tau)$. If $w \equiv -\sqrt{d} \bmod p$, then $bu^2 = (-\sqrt{d} - \tau)(1 + O(p))$. Otherwise if $v(\mu) \geq v(mz^2/d) \geq 1$, then

$$\begin{aligned} bu^2 &= -2\sqrt{d}((mz^2/d - \frac{\mu}{2\sqrt{d}})^2 - (\frac{\sqrt{d}-\tau}{2\sqrt{d}})^2)(1 + O(p)) \\ &= (-\sqrt{d} - \tau)(mz^2/d - \frac{\mu}{2\sqrt{d}})^2(1 + O(p)); \end{aligned}$$

if $v(\mu) < v(mz^2/d)$, then

$$bu^2 = (\sqrt{d} - \tau)(1 + O(p)).$$

Thus $\boxed{p \nmid b, \left(\frac{d}{p}\right) = 1, \left(\frac{b}{p}\right) = \left(\frac{\sqrt{d}-\tau}{p}\right)}$.

(ii) If $v(w) = 1 + 2v(z), v(t) \geq 1 + v(z)$, we may assume $z = 1, v(w) = 1, v(t) \geq 1$. Let $w = pw_1, t = pt_1$, then

$$\begin{aligned} w_1^2 &= d(p^2t_1^4 - \left(\frac{n}{pd}\right)^2), \\ w_1 &\equiv \pm\sqrt{-d}\frac{n}{pd} \pmod{p}, \\ bu^2/p &\equiv (\pm\sqrt{-d} + \mu)\frac{n}{pd} \pmod{p}. \end{aligned}$$

Thus $\boxed{p \mid b, \left(\frac{-d}{p}\right) = 1, \left(\frac{n/b}{p}\right) = \left(\frac{d}{p}\right)\left(\frac{\sqrt{-d}+\mu}{p}\right)}$.

Lemma 3.6. *The curve $\mathcal{M}_b$ defined by* (12) *is locally solvable at*

(1) $p \mid d$ *if and only if*

$$\textit{either} \;\; p \mid b, \; \left(\frac{n/d}{p}\right) = \left(\frac{-n/b}{p}\right) = 1 \;\textit{ or }\; p \nmid b, \; \left(\frac{-n/d}{p}\right) = \left(\frac{-b}{p}\right) = 1;$$

(2) $p \mid \frac{n}{d}$ *if and only if*

$$\textit{either} \;\; p \mid b, \; \left(\frac{-d}{p}\right) = \left(\frac{-2n/b}{p}\right) = 1 \;\textit{ or }\; p \nmid b, \; \left(\frac{d}{p}\right) = \left(\frac{-2b}{p}\right) = 1.$$

Proof. Since d is odd, we may choose $= 1, \tau = (1 + d)/2, \mu = (1 - d)/2$. Then $\sqrt{-d} + \mu = (\sqrt{-d} + 1)^2/2$, $\sqrt{d} - \tau = -(\sqrt{d} - 1)^2/2$. □

3.3. **Proof of main result.** In this subsection, we suppose

$$n = m \text{ or } 2m, m = p_1 \cdots p_k \text{ and } p_i \equiv 3 \pmod 8.$$

Recall for an integer a prime to m, the Jacobi symbol $\left(\dfrac{a}{m}\right) = \prod_{p \mid m} \left(\dfrac{a}{p}\right)$, which is extended to a multiplicative homomorphism from $\{a \in \mathbb{Q}^\times/\mathbb{Q}^{\times 2} : \mathrm{ord}_p(a) \text{ even for } p \mid m\}$ to $\{\pm 1\}$. Set

$$\left[\frac{a}{m}\right] := \frac{1}{2}\Big(1 - \Big(\frac{a}{m}\Big)\Big). \tag{13}$$

The symbol $\left[\dfrac{-}{m}\right]$ is an additive homomorphism from $\{a \in \mathbb{Q}^\times/\mathbb{Q}^{\times 2} : \mathrm{ord}_p(a) \text{ even for } p \mid m\}$ to $\mathbb{F}_2$.

Let A be the $k \times k$ matrix over $\mathbb{F}_2$ with (i,j)-entry $\left[\dfrac{p_j}{p_i}\right]$ $(i \neq j)$ and (i,i)-entry $\left[\dfrac{m/p_i}{p_i}\right]$. For $0 < d \in S^{(\psi)}(E'/\mathbb{Q})$, let $\boldsymbol{d}$ be a vector over $\mathbb{F}_2$ with i-entry 1 if $p_i \mid d$, 0 if $p_i \nmid d$. Then the i-entry of $A\boldsymbol{d}$ is $\left[\dfrac{d}{p_i}\right]$ or $\left[\dfrac{n/d}{p_i}\right]$. Similarly, for $b \in \mathbb{Q}(S, 2), 2 \nmid b > 0$, we have $\boldsymbol{b}$. By Lemma 3.6, we have

$$\begin{aligned}
\text{if } p \mid d,\ \left[\frac{n/d}{p}\right] &= 0, \text{ then } p \mid b,\ \left[\frac{n/b}{p}\right] = \left[\frac{-1}{p}\right] = 1;\\
\text{if } p \mid d,\ \left[\frac{n/d}{p}\right] &= 1, \text{ then } p \nmid b,\ \left[\frac{b}{p}\right] = \left[\frac{-1}{p}\right] = 1;\\
\text{if } p \nmid d,\ \left[\frac{d}{p}\right] &= 0, \text{ then } p \nmid b,\ \left[\frac{b}{p}\right] = \left[\frac{-2}{p}\right] = 0;\\
\text{if } p \nmid d,\ \left[\frac{d}{p}\right] &= 1, \text{ then } p \mid b,\ \left[\frac{n/b}{p}\right] = \left[\frac{-2}{p}\right] = 0.
\end{aligned}$$

If $n = m$, $\boldsymbol{b} = A\boldsymbol{d} + \boldsymbol{d} = (A + I)\boldsymbol{d}$ and $A\boldsymbol{b} = \boldsymbol{d}$. Thus $(A^2 + A + I)\boldsymbol{d} = 0$. Similarly, for $-b, 2b, -2b$, we have $(A^2 + A + I)\boldsymbol{d} = \mathbf{1}, \mathbf{0}, \mathbf{1}$, here $\mathbf{1} = (1, \ldots, 1)^t$. Similarly for $-d$, we have $\boldsymbol{b} = (A + I)\boldsymbol{d} + \mathbf{1}$, but $A\mathbf{1} = \mathbf{0}$, we still have $(A^2 + A + I)\boldsymbol{d} = \mathbf{0}, \mathbf{1}$.

If $n = 2m$, similarly, we have $\boldsymbol{b} = A\boldsymbol{d}$ and $A\boldsymbol{b} = A\boldsymbol{d} + \boldsymbol{d}$. Thus $(A^2 + A + I)\boldsymbol{d} = 0$. Similarly, for $-b, 2b, -2b$, we have $(A^2 + A + I)\boldsymbol{d} = \mathbf{1}, \mathbf{0}, \mathbf{1}$. Similarly for $-d$, we have $\boldsymbol{b} = A\boldsymbol{d} + \mathbf{1}$, but $A\mathbf{1} = \mathbf{0}$, we still have $(A^2 + A + I)\boldsymbol{d} = \mathbf{0}, \mathbf{1}$.

Lemma 3.7. (1) *If* $n = p_1 \cdots p_k$ *and* $p_i \equiv 3 \pmod 8$, *then*

$$\begin{aligned}
S^{(\varphi)}(E/\mathbb{Q}) &= \{1\},\\
S^{(\phi)}(E'/\mathbb{Q}) &= \begin{cases} \{\pm d : 0 < d \mid n, d \equiv 1 \pmod 8\}, & \text{if } 2 \mid k;\\ \{\pm d : 0 < d \mid n\}, & \text{if } 2 \nmid k.\end{cases}
\end{aligned}$$

and $\pm d \in \tilde{S}^{(\psi)}(E'/\mathbb{Q})$ *only if* $(A^2 + A + I)\boldsymbol{d} = \mathbf{0}$ *or* $\mathbf{1}$.

(2) *If* $n = 2p_1 \cdots p_k$ *and* $p_i \equiv 3 \pmod 8$, *then*

$$\begin{aligned}
S^{(\varphi)}(E/\mathbb{Q}) &= \{1\},\\
S^{(\phi)}(E'/\mathbb{Q}) &= \{\pm d : 0 < d \mid n\}.
\end{aligned}$$

and $\pm d, \pm 2d \in \tilde{S}^{(\psi)}(E'/\mathbb{Q})$ *only if* $(A^2 + A + I)\boldsymbol{d} = \mathbf{0}$ *or* $\mathbf{1}$.

Proof of Theorem 1.1. Since $E(\mathbb{Q})_{\mathrm{tor}} \cap \psi E'(\mathbb{Q}) = \{O\}$ and $\#E(\mathbb{Q})_{\mathrm{tor}} = 4$, we have $\#E(\mathbb{Q})/\psi E'(\mathbb{Q}) \geq 4$ and then $\#\tilde{S}^{(\psi)}(E'/\mathbb{Q}) \geq 4$. By Lemma 3.7, if n is odd, $\#\tilde{S}^{(\psi)}(E'/\mathbb{Q}) = 4$; if n is even, $\#\tilde{S}^{(\psi)}(E'/\mathbb{Q}) \leq 8$, but by the existence of the Cassels' skew-symmetric bilinear form on Ш, the $\mathbb{F}_2$-rank of $S^{(\psi)}$ and $\tilde{S}^{(\psi)}$ has same parity, which implies $\#\tilde{S}^{(\psi)}(E'/\mathbb{Q}) = 4$.

Then $\#E(\mathbb{Q})/\psi E'(\mathbb{Q}) = 4$. Since $S^{(\varphi)}(E/\mathbb{Q}) = \{1\}$, $\#E'(\mathbb{Q})/\varphi E(\mathbb{Q}) = 1$. Then

$$\begin{aligned} &\#E(\mathbb{Q})/2E(\mathbb{Q}) = 4,\ \operatorname{rank} E(\mathbb{Q}) = 0,\ \#S^{(2)}(E/\mathbb{Q}) = 4, \\ &Ш(E/\mathbb{Q})[2] = Ш(E/\mathbb{Q})[\varphi] = 0, \\ &Ш(E'/\mathbb{Q})[2] = Ш(E'/\mathbb{Q})[\psi] \cong S^{(\psi)}(E'/\mathbb{Q})/(\mathbb{Z}/2\mathbb{Z})^2. \end{aligned}$$

Hence $Ш(E/\mathbb{Q})[2^\infty] = 0$ and $Ш(E/\mathbb{Q})[2^k\varphi] = 0$. By the exact sequence

$$0 \to Ш(E'/\mathbb{Q})[\psi] \to Ш(E'/\mathbb{Q})[2^k] \to Ш(E/\mathbb{Q})[2^{k-1}\varphi],$$

we have for every $k \in \mathbb{N}_+$,

$$Ш(E'/\mathbb{Q})[2^k] \cong Ш(E'/\mathbb{Q})[\psi],$$

and thus $Ш(E'/\mathbb{Q})[2^\infty] \cong Ш(E'/\mathbb{Q})[\psi] \cong S^{(\psi)}(E'/\mathbb{Q})/(\mathbb{Z}/2\mathbb{Z})^2$. ☐

Proof of Corollary 1.2. In this case, we have

$$A = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ 0 & 0 & 1 & \cdots & 1 \\ 0 & 0 & 1 & \cdots & 1 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \text{ or } \begin{pmatrix} 0 & 1 & 1 & \cdots & 1 \\ 0 & 1 & 1 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix},$$

then $A^2 = A, A^2 + A + I = I$ is invertible. ☐

References

[1] B. Birch, H. P. F. Swinnerton-Dyer. *Notes on ellptic curves (II).* J Reine Angrew Math, 1965, 218:79-108.

[2] J. W. S. Cassels, *Arithmetic on curves of genus 1, (IV) proof of the Hauptvermutung.* J Reine Angrew Math, 1962, 211:95-112.

[3] D. Li, Y. Tian, *On the Birch-Swinnerton-Dyer Conjecture of Elliptic Curves $E_D : y^2 = x^3 - D^2x$*, Acta Mathematica Sinica, 2000, Vol. 16, No. 2, p. 229-236.

[4] Y. Ouyang, S. Zhang, *On non-congruent numbers with* 1 *modulo* 4 *prime factors*, preprint.

[5] J. P. Serre, *A course in Arithmetic*, Springer-Verlag, 1973.

[6] J. H. Silverman, *The Arithmetic of Elliptic Curves*, GTM 106, Springer-Verlag, New York, 1986.

[7] M. Xiong, A. Zaharescu, *Selmer groups and Tate-Shararevich groups for the congruent number problem*, Comment Math Helv, 84(2009), no.1, 21-56.

E-mail address: `zsxqq@mail.ustc.edu.cn`